



## College Counseling Interpretive Report

MMPI®-2

The Minnesota Report™: Adult Clinical System-Revised, 4th Edition

*James N. Butcher, PhD*

Name:	Elton W.
ID Number:	2517
Age:	18
Gender:	Male
Marital Status:	Never Married
Years of Education:	12
Date Assessed:	1/31/14



Copyright © 1989, 1993, 2001, 2005 by the Regents of the University of Minnesota. All rights reserved. Portions reproduced from the MMPI-2 test booklet. Copyright © 1942, 1943 (renewed 1970), 1989 by the Regents of the University of Minnesota. All rights reserved. Portions excerpted from the *MMPI-2 Manual for Administration, Scoring, and Interpretation, Revised Edition*. Copyright © 2001 by the Regents of the University of Minnesota. All rights reserved. Distributed exclusively under license from the University of Minnesota by NCS Pearson, Inc.


**Minnesota Multiphasic Personality Inventory** and **MMPI** are registered trademarks and **The Minnesota Report** is a trademark of the University of Minnesota. **Pearson**, the **PSI logo**, and **PsychCorp** are trademarks in the U.S. and/or other countries of Pearson Education, Inc., or its affiliate(s).

**TRADE SECRET INFORMATION**

Not for release under HIPAA or other data disclosure laws that exempt trade secrets from disclosure.

[ 9.5 / 1 / QG ]

### MMPI-2 VALIDITY PATTERN




Raw Score:	7	8	3	5	4	7	17	28
T Score:	57	57F	45	63	70	65	54	53
Response %:	100	100	100	100	100	100	100	100

Cannot Say (Raw): 0  
 Percent True: 41  
 Percent False: 59

	Raw Score	T Score	Resp. %
S1 - Beliefs in Human Goodness	7	49	100
S2 - Serenity	7	53	100
S3 - Contentment with Life	3	45	100
S4 - Patience/Denial of Irritability	5	54	100
S5 - Denial of Moral Flaws	5	65	100

### MMPI-2 CLINICAL AND SUPPLEMENTARY SCALES PROFILE



Raw Score:	9	24	24	16	15	13	17	13	18	28	22	28	2	7	25	*
K Correction:	9			7			17	17	3							
T Score:	64	62	57	50	30	61	66	56	51	54	53	63	46	48	58	*
Response %:	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	*

Welsh Code: 7+126-38094/:5# L+-K/F:

Profile Elevation: 58.4

\*MDS scores are reported only for clients who indicate that they are married or separated.



## PROFILE VALIDITY

This is a marginally valid MMPI-2 clinical profile because the client may have attempted to present an unrealistically favorable picture of his personal virtue and moral values. He may feel the need to present an image of strong moral character or to deny human frailties. His approach to the MMPI-2 items suggests a rather naive or unsophisticated self-image. Such a pattern could result from reliance on defense mechanisms such as repression or denial. His responses suggest an inflexible life adjustment that may lead to the development of psychological symptoms when stress is present.

## SYMPTOMATIC PATTERNS

Scale Pt was used as the prototype to develop this report. Sensitive and somewhat emotional, the client tends to worry about small matters. He is quite conscientious, something of a perfectionist, rather dissatisfied with himself, and a bit unhappy about his life. In addition, he is concerned about being accepted by others. Individuals with this pattern usually lack confidence in themselves. They are often indecisive, even about everyday matters. Many individuals with this pattern are prone to worry and may become anxious at times without apparent cause. His proneness to anxiety and self-critical attitudes may result in periods of intense stress or tension.

The client has a low Mf score, suggesting that he has a rather limited range of interests and tends to prefer stereotyped masculine activities over literary and artistic pursuits or introspective experiences. He tends to be somewhat competitive and needs to see himself as masculine. He probably prefers to view women in subservient roles. Interpersonally, he is likely to be intolerant and insensitive, and others may find him rather crude, coarse, or narrow-minded. His high endorsement of general anxiety content is likely to be important to understanding his clinical picture.

## PROFILE FREQUENCY

It is usually valuable in MMPI-2 clinical profile interpretation to consider the relative frequency of a given profile pattern in various settings. The client's MMPI-2 high-point clinical scale score (Pt) was found in only 4.9% of the MMPI-2 normative sample of men. Only 3.1% of the sample had Pt as the peak score at or above a T score of 65, and only 1.6% had well-defined Pt spikes.

His MMPI-2 high-point clinical scale score (Pt) was found in 11.1% of the combined sample of college men (Butcher, Graham, Dahlstrom, & Bowman, 1990, sample = 8.5%; Ben-Porath, 1993, sample = 13.0%). Moreover, 7.1% of the students in the combined sample of college men had Pt scale peaks at or above a T score of 65, and 3.7% had well-defined Pt spikes in that range.

## PROFILE STABILITY

The relative elevation of his clinical scale scores suggests that his profile is not as well defined as many other profiles. There could be some shifting of the most prominent scale elevations in the profile code if he were to be retested. The difference between the profile type used to develop the present report

(reflecting the high point on Pt) and the next highest scale in the profile code was 2 points. So, for example, if the client is tested at a later date, his profile might involve more behavioral elements related to elevations on Hs. If so, then on retesting, he might report more physical complaints.

## **INTERPERSONAL RELATIONS**

He tends not to enjoy social activities much. He stands back from participating in groups, where he often feels anxious. He is considered to be hard to get to know and may be somewhat judgmental and perfectionistic, even with close friends. He may be rather critical of the behavior of others. His feelings of inadequacy may impair intimate relationships.

## **DIAGNOSTIC CONSIDERATIONS**

Anxiety is likely to be central in any diagnostic formulation.

## **TREATMENT CONSIDERATIONS**

Individuals with this profile often seek help for their concerns and general unhappiness. Although generally motivated for treatment, they tend to resist psychological interpretations and may rationalize a great deal. It is difficult for them to focus on specific problems. They tend to remain in therapy, but their intellectualization and circular ruminations make progress slow.

**ADDITIONAL SCALES**

	Raw Score	T Score	Resp %
<b>Personality Psychopathology Five (PSY-5) Scales</b>			
Aggressiveness (AGGR)	7	45	100
Psychoticism (PSYC)	5	56	100
Disconstraint (DISC)	13	46	100
Negative Emotionality/Neuroticism (NEGE)	10	51	100
Introversion/Low Positive Emotionality (INTR)	5	37	100
<b>Supplementary Scales</b>			
Anxiety (A)	21	65	100
Repression (R)	16	52	100
Ego Strength (Es)	36	47	100
Dominance (Do)	13	38	100
Social Responsibility (Re)	16	39	100
<b>Harris-Lingoes Subscales</b>			
<b>Depression Subscales</b>			
Subjective Depression (D <sub>1</sub> )	9	56	100
Psychomotor Retardation (D <sub>2</sub> )	6	54	100
Physical Malfunctioning (D <sub>3</sub> )	5	67	100
Mental Dullness (D <sub>4</sub> )	3	53	100
Brooding (D <sub>5</sub> )	1	45	100
<b>Hysteria Subscales</b>			
Denial of Social Anxiety (Hy <sub>1</sub> )	5	56	100
Need for Affection (Hy <sub>2</sub> )	7	51	100
Lassitude-Malaise (Hy <sub>3</sub> )	2	48	100
Somatic Complaints (Hy <sub>4</sub> )	7	72	100
Inhibition of Aggression (Hy <sub>5</sub> )	3	48	100
<b>Psychopathic Deviate Subscales</b>			
Familial Discord (Pd <sub>1</sub> )	1	45	100
Authority Problems (Pd <sub>2</sub> )	2	40	100
Social Imperturbability (Pd <sub>3</sub> )	5	57	100
Social Alienation (Pd <sub>4</sub> )	4	50	100
Self-Alienation (Pd <sub>5</sub> )	3	48	100
<b>Paranoia Subscales</b>			
Persecutory Ideas (Pa <sub>1</sub> )	5	70	100
Poignancy (Pa <sub>2</sub> )	1	41	100
Naivete (Pa <sub>3</sub> )	6	56	100

	Raw Score	T Score	Resp %
<b>Schizophrenia Subscales</b>			
Social Alienation (Sc <sub>1</sub> )	1	43	100
Emotional Alienation (Sc <sub>2</sub> )	1	50	100
Lack of Ego Mastery, Cognitive (Sc <sub>3</sub> )	4	66	100
Lack of Ego Mastery, Conative (Sc <sub>4</sub> )	5	65	100
Lack of Ego Mastery, Defective Inhibition (Sc <sub>5</sub> )	2	54	100
Bizarre Sensory Experiences (Sc <sub>6</sub> )	5	65	100
<b>Hypomania Subscales</b>			
Amorality (Ma <sub>1</sub> )	1	42	100
Psychomotor Acceleration (Ma <sub>2</sub> )	7	58	100
Imperturbability (Ma <sub>3</sub> )	4	53	100
Ego Inflation (Ma <sub>4</sub> )	3	50	100
<b>Social Introversion Subscales (Ben-Porath, Hostetler, Butcher, &amp; Graham)</b>			
Shyness/Self-Consciousness (Si <sub>1</sub> )	5	51	100
Social Avoidance (Si <sub>2</sub> )	0	37	100
Alienation--Self and Others (Si <sub>3</sub> )	9	62	100
<b>Content Component Scales (Ben-Porath &amp; Sherwood)</b>			
<b>Fears Subscales</b>			
Generalized Fearfulness (FRS <sub>1</sub> )	1	53	100
Multiple Fears (FRS <sub>2</sub> )	7	67	100
<b>Depression Subscales</b>			
Lack of Drive (DEP <sub>1</sub> )	1	46	100
Dysphoria (DEP <sub>2</sub> )	0	42	100
Self-Depreciation (DEP <sub>3</sub> )	1	48	100
Suicidal Ideation (DEP <sub>4</sub> )	0	45	100
<b>Health Concerns Subscales</b>			
Gastrointestinal Symptoms (HEA <sub>1</sub> )	2	70	100
Neurological Symptoms (HEA <sub>2</sub> )	2	54	100
General Health Concerns (HEA <sub>3</sub> )	2	56	100
<b>Bizarre Mentation Subscales</b>			
Psychotic Symptomatology (BIZ <sub>1</sub> )	1	54	100
Schizotypal Characteristics (BIZ <sub>2</sub> )	3	60	100
<b>Anger Subscales</b>			
Explosive Behavior (ANG <sub>1</sub> )	0	39	100
Irritability (ANG <sub>2</sub> )	4	56	100
<b>Cynicism Subscales</b>			
Misanthropic Beliefs (CYN <sub>1</sub> )	6	50	100
Interpersonal Suspiciousness (CYN <sub>2</sub> )	6	62	100



	Raw Score	T Score	Resp %
<b>Antisocial Practices Subscales</b>			
Antisocial Attitudes (ASP <sub>1</sub> )	7	52	100
Antisocial Behavior (ASP <sub>2</sub> )	0	38	100
<b>Type A Subscales</b>			
Impatience (TPA <sub>1</sub> )	3	51	100
Competitive Drive (TPA <sub>2</sub> )	5	60	100
<b>Low Self-Esteem Subscales</b>			
Self-Doubt (LSE <sub>1</sub> )	1	44	100
Submissiveness (LSE <sub>2</sub> )	3	62	100
<b>Social Discomfort Subscales</b>			
Introversion (SOD <sub>1</sub> )	0	36	100
Shyness (SOD <sub>2</sub> )	4	58	100
<b>Family Problems Subscales</b>			
Family Discord (FAM <sub>1</sub> )	1	40	100
Familial Alienation (FAM <sub>2</sub> )	1	49	100
<b>Negative Treatment Indicators Subscales</b>			
Low Motivation (TRT <sub>1</sub> )	2	54	100
Inability to Disclose (TRT <sub>2</sub> )	0	37	100

Uniform T scores are used for Hs, D, Hy, Pd, Pa, Pt, Sc, Ma, the content scales, the content component scales, and the PSY-5 scales. The remaining scales and subscales use linear T scores.

## CRITICAL ITEMS

The following critical items have been found to have possible significance in analyzing a client's problem situation. Although these items may serve as a source of hypotheses for further investigation, caution should be used in interpreting individual items because they may have been checked inadvertently.

The percentages of endorsement for each critical item by various reference groups are presented in brackets following the listing of the item. The endorsement percentage labeled "N" is the percentage of the MMPI-2 normative sample of 1,138 men who endorsed the item in the scored direction. Endorsement percentages for the normative sample are reported for all critical items. When available, endorsement percentages for the setting are also reported. The designation "Co" refers to a sample of 1,177 male college students from a combined group of students provided by Butcher, Graham, Dahlstrom, and Bowman (1990) and Ben-Porath (1993).

### Acute Anxiety State (Koss-Butcher Critical Items)

Of the 17 possible items in this section, 5 were endorsed in the scored direction:

- 5. Item Content Omitted. (True) [N = 41; Co = 35]
- 140. Item Content Omitted. (False) [N = 23; Co = 41]
- 172. Item Content Omitted. (True) [N = 9; Co = 15]
- 301. Item Content Omitted. (True) [N = 15; Co = 23]
- 463. Item Content Omitted. (True) [N = 4; Co = 8]



#### Special Note:

The content of the test items is included in the actual reports. To protect the integrity of the test, the item content does not appear in this sample report.

### Depressed Suicidal Ideation (Koss-Butcher Critical Items)

Of the 22 possible items in this section, 3 were endorsed in the scored direction:

- 71. Item Content Omitted. (True) [N = 31; Co = 37]
- 233. Item Content Omitted. (True) [N = 35; Co = 35]
- 273. Item Content Omitted. (True) [N = 16; Co = 20]

### Mental Confusion (Koss-Butcher Critical Items)

Of the 11 possible items in this section, 4 were endorsed in the scored direction:

- 31. Item Content Omitted. (True) [N = 13; Co = 24]

- 299. Item Content Omitted. (True) [N = 15; Co = 25]
- 311. Item Content Omitted. (True) [N = 8; Co = 11]
- 325. Item Content Omitted. (True) [N = 19; Co = 31]



**Special Note:**

The content of the test items is included in the actual reports. To protect the integrity of the test, the item content does not appear in this sample report.

**Persecutory Ideas (Koss-Butcher Critical Items)**

Of the 16 possible items in this section, 4 were endorsed in the scored direction:

- 124. Item Content Omitted. (True) [N = 29; Co = 36]
  
- 251. Item Content Omitted. (True) [N = 24; Co = 43]
- 314. Item Content Omitted. (False) [N = 12; Co = 14]
- 333. Item Content Omitted. (True) [N = 6; Co = 13]

**Antisocial Attitude (Lachar-Wrobel Critical Items)**

Of the 9 possible items in this section, 2 were endorsed in the scored direction:

- 227. Item Content Omitted. (True) [N = 40; Co = 61]
- 254. Item Content Omitted. (True) [N = 24; Co = 32]

**Somatic Symptoms (Lachar-Wrobel Critical Items)**

Of the 23 possible items in this section, 6 were endorsed in the scored direction:

- 47. Item Content Omitted. (False) [N = 19; Co = 23]
- 164. Item Content Omitted. (False) [N = 9; Co = 11]
- 176. Item Content Omitted. (False) [N = 15; Co = 17]
- 224. Item Content Omitted. (False) [N = 18; Co = 14]
- 255. Item Content Omitted. (False) [N = 22; Co = 20]
- 464. Item Content Omitted. (True) [N = 25; Co = 28]

### **Anxiety and Tension (Lachar-Wrobel Critical Items)**

Of the 11 possible items in this section, 5 were endorsed in the scored direction:

- 172. Item Content Omitted. (True) [N = 9; Co = 15]
- 261. Item Content Omitted. (False) [N = 44]
- 299. Item Content Omitted. (True) [N = 15; Co = 25]
- 301. Item Content Omitted. (True) [N = 15; Co = 23]
- 463. Item Content Omitted. (True) [N = 4; Co = 8]



#### **Special Note:**

The content of the test items is included in the actual reports. To protect the integrity of the test, the item content does not appear in this sample report.

### **Sleep Disturbance (Lachar-Wrobel Critical Items)**

Of the 6 possible items in this section, 3 were endorsed in the scored direction:

- 5. Item Content Omitted. (True) [N = 41; Co = 35]
- 140. Item Content Omitted. (False) [N = 23; Co = 41]
- 328. Item Content Omitted. (True) [N = 23]

### **Deviant Thinking and Experience (Lachar-Wrobel Critical Items)**

Of the 10 possible items in this section, 2 were endorsed in the scored direction:

- 122. Item Content Omitted. (True) [N = 80]
- 427. Item Content Omitted. (False) [N = 17]

### **Depression and Worry (Lachar-Wrobel Critical Items)**

Of the 16 possible items in this section, 3 were endorsed in the scored direction:

- 273. Item Content Omitted. (True) [N = 16; Co = 20]
- 339. Item Content Omitted. (True) [N = 37]
- 415. Item Content Omitted. (True) [N = 27]

### Deviant Beliefs (Lachar-Wrobel Critical Items)

Of the 15 possible items in this section, 5 were endorsed in the scored direction:

- 106. Item Content Omitted. (False) [N = 18]
- 314. Item Content Omitted. (False) [N = 12; Co = 14]
- 333. Item Content Omitted. (True) [N = 6; Co = 13]
- 355. Item Content Omitted. (True) [N = 1]
- 466. Item Content Omitted. (True) [N = 32]



#### Special Note:

The content of the test items is included in the actual reports. To protect the integrity of the test, the item content does not appear in this sample report.

### End of Report

---

NOTE: This MMPI-2 interpretation can serve as a useful source of hypotheses about clients. This report is based on objectively derived scale indices and scale interpretations that have been developed in diverse groups of patients. The personality descriptions, inferences, and recommendations contained herein need to be verified by other sources of clinical information because individual clients may not fully match the prototype. The information in this report should only be used by a trained and qualified test interpreter. The report was not designed or intended to be provided directly to clients. The information contained in the report is technical and was developed to aid professional interpretation.

This and previous pages of this report contain trade secrets and are not to be released in response to requests under HIPAA (or any other data disclosure law that exempts trade secret information from release). Further, release in response to litigation discovery demands should be made only in accordance with your profession's ethical guidelines and under an appropriate protective order.

---